



Robotically Assisted Laparoscopic Prostatectomy: Kent and Canterbury Hospital

Information for patients from the Urology Department

This leaflet has been produced to explain **Robotically Assisted Radical Prostatectomy**, as a surgical treatment for prostate cancer. This information supports your consultant and specialist nurse consultations. For more information on the procedure and what will happen when you arrive at hospital, please go to the Prostate Forum web page www.ekhft.nhs.uk/robotic-prostatectomy/

What is Robotic Prostatectomy?

Prostatectomy is the surgical removal of the prostate gland as a treatment for prostate cancer. To do this we now use a surgical robot called the Da Vinci® which enables us to do difficult laparoscopic or keyhole surgery, as opposed to the traditional open surgery. The surgery is performed under a general anaesthetic (you are asleep for the procedure).

The robot is controlled by the surgeon at all times and does not work on its own. The robot is connected to the patient and very small instruments are inserted through small surgical incisions (cuts) in the abdominal wall. The surgeon sits at a separate console and carries out the operation without touching the patient. The surgeon has complete control of the robot at all times.



Kent and Canterbury Hospital was the first hospital in the South East of England, outside of London, to offer this advanced treatment option.



We are a robotic training centre

We are a national training centre in robotic surgery, and run an international fellowship training programme. The fellow works under the supervision of consultants at all times, and may perform part or all of your surgery.

Is the surgeon able to do the same operation as an open prostatectomy with the robot?

Yes. In terms of cancer treatment it is regarded as the same as open surgery and does not compromise cancer control.

What are the advantages of robotic surgery?

This type of surgery offers many advantages for both you and your surgeon.


Surgical advantages

- A highly magnified high definition view of your operation
- A complete 3D view of your prostate when operating
- Much finer more delicate dissection
- Far more precise surgery
- It eliminates tremor.

Patient advantages

- Lower complication rates
- Lower infection rates
- Less pain
- Quicker recovery
- Shorter hospital stay
- Shorter catheter time
- Less blood loss and a lower rate of needing a blood transfusion (1% or one in every 100 patients)
- Earlier return to work/normal daily activities
- Improved cosmetic result.

Comparisons between incision sites



For more information, please visit the Trust's Prostate Forum web page www.ekhuft.nhs.uk/robotic-prostatectomy/

Are there any alternatives to this treatment?

Yes. These should have been discussed with you by your urologist or oncologist. Depending on your age, PSA blood test, type of disease, and other health issues, radiotherapy, brachytherapy, surveillance, or hormone therapy are all treatment options. However this type of surgery is only chosen if we feel it is the most appropriate treatment option for you, and you are happy to proceed.

Are there any complications of this surgery?

Yes. All surgery has the potential for complications, however robotic surgery minimises these risks. These should be discussed with you by your surgeon before signing the consent form.

Short term common risks

- Bruising in the scrotum and penis and/or around the wounds – wear supportive underwear following surgery for two to three weeks (very common).
- More severe bruising across the abdomen, this happens rarely and resolves over the first couple of weeks. It does not necessarily mean there is a problem with the operation (uncommon).
- Bladder spasms/pain with catheter (common).
- Catheter bypassing – urine may leak down the side of the catheter causing minor leakage, this is common and not a concern.
- Constipation – this is important to avoid and you will be given medication for this (uncommon).
- Blood in the urine (common).
- Urinary or wound infection (uncommon).
- Facial swelling lasting a few hours (common).
- Shoulder pain lasting a few hours (very common).
- Rectal discomfort or a desire to open the bowels which can last for the first hour or two after the operation (common).
- A feeling of abdominal bloating or gas inside the abdomen, this can last for a few days after discharge and settles once the bowels are opened regularly (very common).

Short term rare complications

- Bleeding needing transfusion during or after the operation (two in 100 patients).
- Return to theatre for complications such as bleeding (one in 200 patients).
- Damage to structures around the prostate for example the bowel/rectum/bladder/blood vessels or ureters (tubes between kidney and bladder). Also potential damage to nerve supply to muscles in the leg can occur rarely (less than one in every 100 patients).
- Urine leak, this is where urine leaks from the join between the bladder and waterpipe. It rarely needs reoperation and just heals up over a few days needing a longer stay in hospital (one in 200 patients).
- Risk of clots in the legs (DVT or deep vein thrombosis), or clots on the lungs (PE or pulmonary embolus), again around (one in 200 patients). You will have injections to reduce this problem. Keeping well hydrated and early mobilisation/walking helps prevent this problem.
- Medical complications such as chest infections or heart attack, again this is rare and is less than one in 200 patients.
- Conversion to an open operation – again this is rare (0%).
- Readmission to hospital (four in 100 patients).
- Urinary retention after catheter removal (three in every 200 patients).
- Port site hernias (bowel protruding through wound) (one in 100 patients).
- Equipment failure where the operation would be cancelled and rescheduled is rare.
- This is major pelvic surgery and as a result there is a risk of death but this is less than one in every 1000 patients.

Removal of lymph nodes

In selected patients with higher grade or higher volume disease, removal of the lymph nodes within the pelvis may be discussed with you. The reason for this is to find out if there has been spread of the prostate cancer to these lymph nodes that is not detectable on scans.

Most patients will not notice any side effects from the removal of the nodes. A small percentage of patients may experience complications related to this part of the operation. These include:

- Lymphoceles – these are collections of fluid in the pelvis. They may cause pain or discomfort which may occasionally need draining (five in every 100 patients).
- Lymph leak – this is where fluid from the lymphatic system is produced that needs drainage from the abdomen, this needs continued drainage at home.
- Lymphoedema or swelling of the leg – if this happens patients are referred to the community lymphoedema team. It can last for nine to 12 months and in rare cases is permanent (one in 100 patients).
- Damage to pelvic nerves or blood vessels can affect the nerve on the inner thigh, which can cause difficulty when crossing your legs (very rare).
- Pelvic Haematomas is a solid swelling of blood in the tissues outside the blood vessels.

Most of the above problems would be identified by your medical team before your discharge from hospital or at your follow-up appointments. If you are at all concerned please use the contact numbers listed on page 8 or printed on your discharge leaflet. You will also be contacted by phone by our specialist nurses following discharge; you can raise any concerns here.

For more information, please visit the Trust's Prostate Forum web page www.ekhuft.nhs.uk/robotic-prostatectomy/

Long term complications

- **Incontinence** - continence can be affected by this treatment. However, long term (longer than a year) incontinence that needs further surgery happens in two to three in every 100 patients. The majority of patients will have some incontinence when their catheter is removed and may need to use incontinence pads for a period of time. With robotic surgery continence recovers quicker.

It is important to do your pelvic floor exercises before, during, and after your recovery. You will be given written information regarding pelvic floor exercises. The community physiotherapists will help in this area, and an appointment will automatically be made for you.

Patients who already have pre-existing stress urinary incontinence and/or urge incontinence are at a higher risk of needing pads long term after surgery.

The table below outlines continence recovery. This is taken from our own audited results.

Time since surgery	% of patients that are pad free
one week	29 to 30%
six weeks	45 to 50%
12 weeks	65 to 75%
26 weeks	75 to 85%
52 weeks	90 to 95%

- **Erectile dysfunction** - erections will be affected by this surgery. This is more likely in older age groups and if you have weak erections before the surgery. In patients who want to regain their erections there may be the possibility that we can offer nerve sparing technique. This is part of the operation where we protect the nerves that go down to the penis, these run very close to the prostate and can be gently peeled off. We do not offer this in patients with more advanced disease; your surgeon will discuss with you if appropriate. Even with nerve spare surgery, erections are not guaranteed. You will be offered a daily tablet to help with recovery. Spontaneous recovery that is going to happen can take up to two years, although we have dedicated clinics for erectile dysfunction that we can refer you to after your surgery.
- Some patients will notice a degree of **penile shortening**. In rare cases this can be severe and permanent.
- **Urethral/Anastomotic strictures** (scar tissue that forms in the water pipe).
- Patients may notice a slowing of urinary flow or struggling to pass urine. It can occur soon or many months after the operation. Anastomotic strictures have occurred in one in every 200 of our patients. Sometimes it needs a minor operation to treat it.

For more information, please visit the Trust's Prostate Forum web page www.ekhft.nhs.uk/robotic-prostatectomy/

Will I lose my desire/libido after surgery?

No. The operation should not affect your libido. It will however affect your fertility. You will have no ejaculate following your operation and in some men they notice a small degree of penile shortening.

When can I start sexual relations?

There is no restriction on resuming sexual relations after the catheter is removed, once you and your partner are happy to do so.

Can the robot breakdown?

Potentially, although this very rarely happens (less than one in every 200 operations). If this happens either your operation can be rescheduled or it can be done as an open operation. Over 4000 robots are being used worldwide and this has only been reported in a handful of cases. We have had to cancel operations on the day of surgery due to equipment failure when testing before surgery, but then rebooked patients onto the next available surgical list.

What happens now?

Once we have decided on robotic surgery:

- You will be contacted with a date for your operation
- You will be asked to attend the preassessment clinic at Kent and Canterbury Hospital where routine bloods tests and an ECG (heart tracing) are taken, and an assessment of your fitness for surgery is made
- You and your wife/partner will be invited to a 'Patients Forum'; this is a group that meets once a month and discusses issues surrounding the surgery, such as continence, physio, and erectile function. The Prostatectomy Education Forum has now been recorded and the videos are available to view via the Trust web site www.ekhuft.nhs.uk/robotic-prostatectomy/
- You will be admitted to the hospital (Clarke Ward, Kent and Canterbury Hospital) on the morning of your operation. You will meet your surgeon, anaesthetist, and nursing staff
- You will be asked to sign the consent form. Remember you can withdraw your consent for treatment at any time
- You will be given an enema before your operation, to make sure your bowels are clear.

How long does the operation take?

Between two to three hours. It is done under general anaesthetic (you are asleep for the operation). You will wake up in the recovery bay before being transferred back to the ward.

For more information, please visit the Trust's Prostate Forum web page www.ekhuft.nhs.uk/robotic-prostatectomy/

Will I be in pain?

Robotic surgery is less painful than open surgery. Before you wake up the anaesthetist will give you pain relief and we use local anaesthetic (numbing solution) in your wounds and bladder. If you are in pain then please tell the nursing staff so they can give you pain relief. A lot of patients wake up feeling as though they have a full bladder or get bladder spasms, this is because of the catheter and the surgery around your bladder, not because your bladder is full. It usually settles but if not can be helped with medication.

The anaesthetist may consider a spinal injection to help with pain control before your surgery.

How long will I be in hospital?

One day. The majority of patients are discharged the day after their operation. Along with the catheter, you will wake up with a drain tube which is removed before your discharge from hospital. You will also be taught how to give yourself a small injection, each day for four weeks, to help prevent clots forming in the veins in your legs.

Preventing blood clots (DVT or deep vein thrombosis)

To prevent clots you will be taught to give yourself an injection into your stomach for 28 days. You will be taught how to do this on the ward before you leave hospital.

You will also be given surgical stockings to wear for a time after surgery.

Will I have help once discharged?

Yes.

- Before leaving hospital you will be given instructions on how to manage your catheter and wound dressings
- Absorbable stitches are used so no stitches need to be removed
- You will be given painkillers to take as and when you need them, tablets to stop gastritis (stomach irritation) to be taken for the first week, and medication to prevent constipation
- For men whom erectile function is important a medication will be offered, and this should be started once the catheter is removed
- If needed the community nurse is available (contact details are available via your GP)
- The Community Continence Team are available and you can refer to them for support
- The ward can be called if there are any problems.

Please see page 10 for useful contact numbers.

Are there any restrictions once I go home?

To start with exercise should be gentle but gradually built up on a daily basis as symptoms allow. Do not stay in bed all the time. You should take a full and balanced high fibre diet and drink plenty of fluids, especially in the first week. Do not drink alcohol for one week after your surgery. Energy levels can vary from day to day following surgery.

For more information, please visit the Trust's Prostate Forum web page www.ekhuft.nhs.uk/robotic-prostatectomy/

How long will I need a catheter for?

The catheter is a very important part of your recovery. It protects the join between your bladder and your urethra (water pipe) in the early stages of the healing process. We like to leave the catheter in place for as short a time as possible, between seven to 10 days after your operation. Removal of the catheter is very straight forward and only takes a matter of seconds; this is done in the Urology department. You will be given a date for catheter removal before you are discharged from hospital. Occasionally your surgeon may make a decision to leave the catheter in for longer, but you would be told of this decision and the reasons why.

Once the catheter is removed please continue doing your pelvic floor exercises regularly.

Remember, that to start with, most patients have some leakage once their catheter is removed and you may need to use pads for a period of time. This is where the Kent Continence Service will help you.

When do I get my results?

It takes our pathologists about two weeks to process and examine your prostate. If you wish to receive your results before your six week follow-up please ask your prostate specialist nurse when you are contacted after discharge.

Will I need any other forms of treatment?

Most patients do not need any other forms of treatment. Some patients with more advanced disease may be offered radiotherapy after their surgery, either immediately or later if there is a rise in the PSA, others may be given a long course of hormones. This will be discussed with you by your surgeon. Further appointments with the oncologists may be needed in this situation.

What follow-up will I receive?

After the initial pathology results, you will be seen by your surgeon six weeks after your surgery, then at three monthly intervals to check your PSA levels. If all is well, then time between appointments can be lengthened. You may also have appointments with the physiotherapist to make sure your pelvic floor muscles are recovering. Erectile dysfunction will be discussed at a specialist clinic, if this is important to you.

When can I drive?

This varies between patients. You will only be fit to drive if you can do an emergency stop without it causing pain. Please wait at least one week before driving again.

For more information, please visit the Trust's Prostate Forum web page www.ekhuft.nhs.uk/robotic-prostatectomy/

When can I return to work?

Again this varies between patients. You can return to work when you feel able. Some patients are back to work after one week, others take three to four weeks. It is important only to do light duties to start with and to do no heavy lifting for four weeks.

We have found that after robotic surgery patients return to work quicker, on average at three weeks compared to 11 weeks after open surgery.

When can I fly?

You can fly on short haul flights after two weeks, and long haul flights after four weeks. These are recommendations, if you have any concerns please delay your travel. You must also check with your carrier and travel insurance company.

What can we do to improve our service?

We are constantly trying to improve the high quality care we provide. We continually audit and assess our results. We will ask you to fill out questionnaires on your continence, erectile dysfunction, and overall quality of life. We would like you to fill these out before your operation and at all your follow-up appointments. We also want you to record the first day you manage without pads and when you return to work. Please let us know these dates at your follow-up appointment or by email.

All this information is handled sensitively. Please feel free to add any other comments, whether good or bad. All these comments will be taken seriously and will be used to improve the experience for future patients.

Where can I get further information about the operation?

- da Vinci Prostatectomy www.davinciprostatectomy.com
- Prostate Cancer UK www.prostatecanceruk.org/
- CancerHelp UK www.cancerhelp.org.uk
- Patient.co.uk www.patient.co.uk
- Macmillan Cancer Support www.macmillan.org.uk

For more information, please visit the Trust's Prostate Forum web page www.ekhft.nhs.uk/robotic-prostatectomy/

Useful contact numbers

- **Macmillan Urology Cancer Nurse Specialists**
Morna Jones / David Stafford / Janey Blofield / Emma Chamberlain
Telephone: 01227 868666
Email: ekhufft.uro-onccns@nhs.net
- **Macmillan Urology Support Worker**
Robyn Dunn and Marisa Smith
Email: ekhufft.urocancersupportworkers@nhs.net
- **Clarke Ward**
Kent and Canterbury Hospital
Telephone: 01227 783103
- **Kent Continence Service**
Telephone: 0300 7900310
- **The Prostate Cancer Charity**
Email: info@prostate-cancer.org.uk
- **Macmillan Cancer Support**
Free helpline: 0808 808 00 00
Practical, financial, and emotional support for people with cancer, their family and friends.
Macmillan also has further information on sex, relationships, and cancer.
- **The Bladder and Bowel Foundation (B&BF)**
Nurse Helpline: 0845 345 0165
- **Sexual Dysfunction Association**
For support and information on erectile dysfunction
Helpline: 0870 774 3571

This leaflet has been produced with and for patients

If you would like this information in **another language, audio, Braille, Easy Read, or large print** please ask a member of staff.

Any complaints, comments, concerns, or compliments please speak to your doctor or nurse, or contact the Patient Advice and Liaison Service (PALS) on 01227 783145, or email ekh-tr.pals@nhs.net

Further patient leaflets are available via the East Kent Hospitals web site www.ekhufft.nhs.uk/patientinformation